


Hermetically  
Sealed in Foil for  
Freshness


# MayFord

## HERBS KRUIE

### ROSEMARY (*Rosmarinus officinalis*)



LOCATION  
LIGGING  
LOCALIZAÇÃO




DIRECT / DIREK / DIREKTAMENTE




WINTER WINTER	SPRING LENTE	SUMMER SOMER	AUTUMN HERFS
INVERNO	PRIMAVERA	VERÃO	OUTONO


SOWING TIME • SAAITYD • ÉPOCA DA SEMEITEIRA




SOWING DEPTH  
SAAIDIEPTE  
PROFUNDIDADE  
DE SEMEADURA



PRESS DOWN  
DRUK VAS  
COMPRIMER



KEEP MOIST  
HOU KLAM  
MANTER ÚMIDO



THIN OUT AT  
DUN UIT TEEN  
RALEAR



THIN OUT TO  
DUN UIT NA  
RALEAR PARA




KEEP MOIST  
HOU KLAM  
MANTER ÚMIDO



FINAL HEIGHT  
FINALE HOOGTE  
ALTURA FINAL



DAYS TO HARVEST  
DAE TOT OES  
DIAS AT É COLHEITA



Slow and erratic germinator. Soaking in water for 2 days before sowing may enhance germination which can take as long as 3 to 4 weeks. Consistent moisture levels are important during germination. Rosemary attracts bees, butterflies and birds. Do not plant near Basil.



30MSP00 - 00632



## SAKATA

SAKATA SEED SOUTHERN AFRICA (Pty) Ltd  
P.O. BOX 160, LANSERIA 1748 SOUTH AFRICA  
[www.mayford.co.za](http://www.mayford.co.za)



Not for food,  
feed or processing

