

LOCATION / LIGGING LOCALIZAÇÃO



FROST FREE AREAS
RYPVRYGEBIEDE
ÁREAS ISENTAS DE GEADA

FROST AREAS / RYPGEBIEDE
ÁREAS GEADAS

WINTER
WINTER



INVERNO

SPRING
LENTE



PRIMAVERA

SUMMER
SOMER




VERÃO

AUTUMN
HERFS



OUTONO

SOWING TIME • SAAITYD • ÉPOCA DA SEMEITEIRA




Super Phosphate Superfostaat Superfostato		30 g / m ²
Lawn Fertiliser Grasperk Kunsmis Fertilizante para relva		30 g / m ²
OR		
Organic Rootbuilder Fertiliser		100 g / m ²



RAKE LEVEL
HARK
GELYK
NIVELAR
COM
ANCINHO




MIX WITH SAND
MENG MET SAND
MISTURAR COM AREIA



SOWING DEPTH
SAAIDIEPTE
APROFUNDIDADE DE SEMEITEIRA


5 - 10 mm



ROLL or TRAMPLE - ROL of TRAP VAS
COM ROLO ou PISAR




KEEP MOIST - AVOID PUDDLES
HOU KLAM - VERMY WATERPOELE
MANTER HÚMIDO - EVITAR POÇAS



Lawn Fertiliser
Grasperk Kunsmis
Fertilizante para relva

30 g / m²
4 x PER YEAR / JAAR / POR ANO

BERMUDA (Premier Blend Cynodon dactylon) is a medium fine textured creeping grass with excellent drought, heat and wear tolerance. It becomes dormant (goes brown) when exposed to frost.



MOW AT
SNY TEEN
CORTAR AOS
2 - 3 cm

CUTTING HEIGHT / SNY HOOGTE / ALTURA DE CORTE

Despite this container serving to prolong the life of the seed, it should not be exposed to unduly high temperatures. Once the container has been torn open, deterioration of the seed will ensue at the usual rate.



6 002788 004878 >



30MSP03 - 00397



SAKATA SEED SOUTHERN AFRICA (Pty) Ltd.
POSTNET SUITE 591, NORTHRIDING 2163
www.mayford.co.za

